

Multi-quintessence

R.A.C. Correa* and A. de Souza Dutra†

UNESP, Universidade Estadual Paulista, 12516-410, Guaratinguetá, SP, Brazil

P.H.R.S. Moraes‡

ITA - Instituto Tecnológico de Aeronáutica - Departamento de Física,
12228-900, São José dos Campos, São Paulo, Brazil

In this work we derive and analyse cosmological scenarios coming from multi-component scalar field models. We consider a direct sum of a sine-Gordon and a Z_2 model and a combination of those with a BNRT model. All the computations are made analytically and some parameters restriction is put forward in order to get in touch with realistic complete cosmological scenarios.

PACS numbers:

Keywords: cosmology, scalar fields, first-order formalism

I. INTRODUCTION

In standard cosmology model (SM), the quantum vacuum energy is the responsible for the cosmic acceleration observationally predicted [1, 2]. It enters the field equations of General Relativity in the form of a cosmological constant. However, there is a huge discrepancy between the theoretically predicted value for the vacuum quantum energy obtained via Particle Physics [3] and via cosmological observations [4]. Such a discrepancy yields the consideration of alternative gravity theories, from which healthy cosmological models can be derived. Among the alternatives we quote $f(R)$ and $f(R, T)$ theories [5–14].

Another possibility comes from scalar field models, named quintessence, in which it is assumed that the dynamics of the Universe is governed by some scalar field ϕ . Depending on the form of the scalar field time evolution and on its potential, it is possible to describe the accelerated phases of the Universe dynamics, namely dark energy and inflationary eras [15, 16]. Quintessence models with two scalar fields driving the Universe dynamics have also generated well-behaved cosmological models [17–20].

In the present work we derive cosmological scenarios from N-field quintessence models, which we call multi-quintessence models. In order to obtain the scalar field solutions we apply the first order formalism [20, 21]. The solutions for the cosmological parameters are compared with observations and we show that for some values of the free parameters of the models, it is possible to draw a complete cosmological scenario.

The paper is organized as follows: in Section II we describe the first order formalism. In Section III we obtain our scalar field solutions and in Section IV we construct the cosmological parameters of the models. In Section V we revisit our main results and present a deep discussion about the physical properties of them.

II. A BRIEF REVIEW OF THE FIRST-ORDER FORMALISM

In this section we show a brief overview about the first-order formalism for the coupling of scalar fields with gravity [20, 21]. Let us begin by writing the action in four-dimensional gravity which is coupled to a set of real scalar fields ϕ_i , $i = 1, 2, \dots, N$, in the following form

$$S = \int d^4x \sqrt{|g|} \left[-\frac{1}{4}R + \frac{1}{2}g_{ab}\nabla^a\phi_i\nabla^b\phi_i - V(\phi_i) \right]. \quad (1)$$

We are using $c = 4\pi G = 1$ and $g = \det(g_{ab})$, with field, space and time variables being dimensionless and R stands for the Ricci scalar. Furthermore, we are adopting the signature of the metric as $(+ - - -)$. Moreover, as usual, $V(\phi_i) \equiv V(\phi_1, \phi_2, \dots, \phi_N)$ is the potential that describes the theory in terms of a finite arbitrary number of scalar fields.

We will study the case where the metric is the Friedmann-Robertson-Walker one, responsible for a homogeneous and isotropic expanding universe,

$$ds^2 = dt^2 - a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2 (d\theta^2 + \sin^2\theta d\varphi^2) \right], \quad (2)$$

with k being the curvature parameter, which will be taken as zero, in accord with observations [4]. From this metric, the equations relating the Hubble and the expanding factors are

$$H^2 = \frac{2}{3}\rho - \frac{k}{a^2}; \quad H \equiv \frac{\dot{a}}{a} \quad (3)$$

and

$$\frac{\ddot{a}}{a} = -\frac{1}{3}(\rho + 3p); \quad \bar{q} = \frac{\ddot{a}}{\dot{a}^2} = 1 + \frac{\dot{H}}{H^2}, \quad (4)$$

with dots representing time derivatives and ρ and p being the energy density and pressure of the system, respectively. Furthermore, it is very useful to work with the equation of state (EoS) parameter

$$\omega = \frac{p}{\rho}. \quad (5)$$

*Electronic address: fis04132@gmail.com

†Electronic address: dutra@feg.unesp.br

‡Electronic address: Moraes.phrs@gmail.com

Let us assume that the fields ϕ_i also depend only on t , that is $\phi_i = \phi_i(t)$. Thus, from action (1) we can obtain the corresponding equations of motion for the scalar fields

$$\ddot{\phi}_i + 3H \dot{\phi}_i + V_{\phi_i} = 0, \quad (6)$$

with $V_{\phi_i} \equiv dV/d\phi_i$.

Considering that the potential is given by

$$V(\phi_i) = \frac{3}{2}W(\phi_1, \phi_2, \dots, \phi_N)^2 - \frac{1}{2} \sum_{i=1}^N W_{\phi_i}^2, \quad (7)$$

with $W(\phi_i)$ being a superpotential and $W_{\phi_i} \equiv dW/d\phi_i$, one can perceive that even if we use a superpotential which is a direct sum of one scalar field functions, the resulting field potential is non-trivial as it could be seen in braneworld scenarios [22]. The energy density and pressure in this formalism are written as

$$\rho = \sum_{j=1}^N \frac{\dot{\phi}_j^2}{3} + \frac{3}{2}W(\phi_1, \phi_2, \dots, \phi_N)^2 - \frac{1}{2} \sum_{i=1}^N W_{\phi_i}^2, \quad (8)$$

$$p = \sum_{j=1}^N \frac{\dot{\phi}_j^2}{3} - \frac{3}{2}W(\phi_1, \phi_2, \dots, \phi_N)^2 + \frac{1}{2} \sum_{i=1}^N W_{\phi_i}^2. \quad (9)$$

Furthermore, in terms of the superpotential, the differential equations of motion can be cast in the first order form when we use the identification $H = -W(\phi_1, \phi_2, \dots, \phi_N)$, and are written as

$$\dot{\phi}_i = W_{\phi_i}, \quad i = 1, \dots, N. \quad (10)$$

From this approach, it was shown in Ref.[22] that some braneworld scenarios allow to find interesting features, such as split brane mechanism and asymmetric warp factor shape, which is essential to address the hierarchy problem in the context of thick brane scenarios. As demonstrated by the authors, the method is not restricted to the combination of one-field nonlinear superpotentials. In fact one could combine two or more superpotentials where two or more coupled scalar fields appear non trivially.

In the next section, we will apply the approach described above to study cosmological scenarios coming from multi-component scalar field models.

III. SCALAR FIELD SOLUTIONS

Now we go further in the approach, by presenting simple, non trivial and analytical examples.

We will start by considering a superpotential which is a direct sum of a sine-Gordon [23] and a Z_2 model [24], as:

$$W(\phi_1, \phi_2) \equiv \lambda_1 \left(\phi_1 - \frac{\phi_1^3}{3} \right) + \lambda_2 \sin \phi_2 + \alpha_1. \quad (11)$$

where α_1 is an arbitrary constant.

At this point it is worth to emphasize that the Z_2 model is the most fundamental example where we can find the so-called Z_2 kink. In this context, the Lagrangian is invariant under the transformation $\phi \rightarrow -\phi$ and hence there is a reflectional Z_2 symmetry. On the other hand, the sine-Gordon model is invariant under $\phi \rightarrow \phi + 2\pi n$, where n is any integer, and thus possesses Z symmetry. The motivation to study this first example, where we have a mixture of sine-Gordon and Z_2 model, comes from the fact that these models are found in many areas of Physics, including condensed matter physics, field theory, cosmology, among many others. For instance, as it is well known, whenever we have a potential with two or more degenerate minima, one can find different vacua at different portions of the space. Thus, one can find domain walls connecting such regions. In this framework, since the Universe is expanding and cooling, cosmic phase transitions can occur and domain walls can form [24].

Thus, using Eq.(10), we obtain the following differential equations

$$\dot{\phi}_1 = \lambda_1 (1 - \phi_1^2), \quad (12)$$

$$\dot{\phi}_2 = \lambda_2 \cos \phi_2. \quad (13)$$

The solutions of Eqs.(12) and (13) are, respectively, given by

$$\phi_1(t) = \tanh(\lambda_1 t), \quad (14)$$

$$\phi_2(t) = \arctan[\sinh(\lambda_2 t)]. \quad (15)$$

A second example can be represented as a combination of Z_2 , sine-Gordon and the so-called BNRT models [25]. In this case, the superpotential can be written in the following form

$$W(\phi_1, \phi_2, \phi_3, \phi_4) = \lambda_1 \left(\phi_1 - \frac{\phi_1^3}{3} \right) + \lambda_2 \sin \phi_2 - \lambda_3 \phi_3 + \frac{\lambda_3}{3} \phi_3^3 + \mu_3 \phi_3 \phi_4^2 + \alpha_2. \quad (16)$$

where α_2 is an arbitrary constant.

Let us remark that the BNRT model [25] is used for modeling a great number of systems [26–30]. This model admits a variety of kink-like solutions and it has been shown to give rise to degenerate and critical branes [31], traveling solitons [32, 33] and BPS and non-BPS defects [34]. Moreover, using the so-called configurational entropy [35], it was shown in Ref.[36] that the model has a rich structure, which allows the generation of double-kink configurations.

From Eq.(16), we have the following first-order equations

$$\dot{\phi}_1 = \lambda_1 (1 - \phi_1^2), \quad (17)$$

$$\dot{\phi}_2 = \lambda_2 \cos \phi_2, \quad (18)$$

$$\dot{\phi}_3 = -\lambda_3 (1 - \phi_3^2) + \mu_3 \phi_4^2, \quad (19)$$

$$\dot{\phi}_4 = 2\mu_3 \phi_3 \phi_4. \quad (20)$$

We can see that Eqs.(17) and (18) also have Eqs.(14) and (15) as solutions, respectively. On the other hand, Eqs.(19) and (20) are coupled. In this case, it has been found in Ref.[37] a class of analytical solutions for this system, which was named Degenerate Bloch Walls (DBW). It was shown that there are two situations with exact classical solutions. The first one is given by

$$\phi_3^{(1)}(t) = \frac{(\sqrt{c_0^2 - 4}) \sinh(2\mu_3 t)}{(\sqrt{c_0^2 - 4}) \cosh(2\mu_3 t) - c_0}, \quad (21)$$

$$\phi_4^{(1)}(t) = \frac{2}{(\sqrt{c_0^2 - 4}) \cosh(2\mu_3 t) - c_0}, \quad (22)$$

where $c_0 < -2$ and $\lambda_3 = \mu_3$.

For the second class of solutions, $c_0 < 1/16$ and $\lambda_3 = 4\mu_3$. In this case, we have

$$\phi_3^{(2)}(t) = \frac{(\sqrt{1 - 16c_0}) \sinh(4\mu_3 t)}{(\sqrt{1 - 16c_0}) \cosh(4\mu_3 t) + 1}, \quad (23)$$

$$\phi_4^{(2)}(t) = -\frac{2}{(\sqrt{1 - 16c_0}) \cosh(4\mu_3 t) + 1}. \quad (24)$$

In Figs.1 and 2 below we show some typical profiles of the DBW solutions. Note that the two-kink solution $\phi_3^{(1)}$ arises only for values of c_0 close to the critical value, $c_0^{crit} = -2$. For the same values of c_0 , the corresponding lump-like solutions for $\phi_4^{(1)}$ exhibit a flat top, which disappears as one moves away from c_0^{crit} . A remarkable feature of these solutions is that their BPS energies [23] are degenerate with respect to c_0 , where for any value of c_0 , the energy is given by $E_{BPS} = 4\lambda/3$.




FIG. 1: Profiles of the DBW solutions $\phi_3^{(1)}(t)$ with $\mu = 1$.

IV. COSMOLOGICAL PARAMETERS SOLUTIONS

Below we plot the cosmological solutions for $H(t)$, $\omega(t)$ and $\bar{q}(t)$ for both models described above, with superpo-




FIG. 2: Profiles of the DBW solutions $\phi_4^{(1)}(t)$ with $\mu = 1$.

tential (11) representing what we will call *Model I* and superpotential (16) representing *Model II*.




FIG. 3: Hubble factor for different values of the c_0 parameter with $\lambda_1 = 2.70$, $\lambda_2 = 3.23$, and $\alpha_1 = -5.9$. The (blue) dotted and (red) thin lines stands for $\lambda_3 = \mu_3 = 0.9$.

In the next section, we will interpret Figs.3-5 and point the features that corroborate to the fact that for some chosen values of the free parameters, some of them are capable of describing a complete cosmological scenario, i.e., one that includes the inflationary, radiation, matter and dark energy dynamical eras [38, 39], as well as the transitions among these stages, in a continuous form.

V. DISCUSSION

The main cosmological puzzle nowadays is to predict in a theoretical level the observed cosmic acceleration. Whether this phenomenon comes from extra degrees of freedom in the geometrical description [8, 40, 41] or material content [9, 14, 42–44] of the Universe is still an open question.

In SM, the cosmic acceleration is considered as due to the presence of the cosmological constant in the Ein-




FIG. 4: Behavior of the equation of state with $\lambda_1 = 2.70$, $\lambda_2 = 3.23$, and $\alpha_1 = -5.9$. The (blue) dotted and (red) thin lines stands for $\lambda_3 = \mu_3 = 0.9$.




FIG. 5: Acceleration parameter evolution for different values of the c_0 parameter with $\lambda_1 = 2.70$, $\lambda_2 = 3.23$, and $\alpha_1 = -5.9$. The (blue) dotted and (red) thin lines stands for $\lambda_3 = \mu_3 = 0.9$.

stein's field equations. As quoted in the Introduction, such a consideration yields the cosmological constant problem. In fact, there is a number of shortcomings or inconsistencies surrounding SM, as one can check in [45, 46], for instance.

Another shortcoming, barely explored in the literature, surrounding the SM is that it cannot describe the whole Universe dynamics continuously. We know that since the Big-Bang the Universe is expanding, however, the dynamical features of this expansion are not the same since then. Right after the Big-Bang, the Universe passed through an accelerated regime of expansion, named inflation (or inflationary era) [47, 48]. The inflaton, which is named the field responsible for bringing such an abrupt dynamics to the Universe decays rapidly in radiation. After radiation, (dark and barionic) matter and dark energy, subsequently, dominate the Universe dynamics [38]. To say that SM is unable of describing the dynamics of the Universe in a continuous form means that the cosmological parameters of the model, such as the EoS and

acceleration parameters, need to be analysed separately for each of the stages mentioned above.

In other words, in SM, one is unable to find an analytical solution, say, for the EoS parameter, which is able to describe the inflationary, radiation, matter and dark energy eras, and the transitions among them in a continuous and self-consistent analytical form. Rather, when solving the standard Friedmann equations, it is admissible to know the EoS of the dominant component of the Universe dynamics at a specific time, be the inflation, radiation, matter or dark energy. Let us clarify this statement by writing the (standard) Friedmann equations and the energy-momentum tensor conservation equation of SM (for $k = 0$):

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{2}{3}\rho, \quad (25)$$

$$\frac{\ddot{a}}{a} + \frac{1}{2}\left(\frac{\dot{a}}{a}\right)^2 = -\omega\rho, \quad (26)$$

$$\dot{\rho} + 3\frac{\dot{a}}{a}(1 + \omega)\rho = 0, \quad (27)$$

in which it was already assumed the EoS $p = \omega\rho$. We can see from (25)-(27) a set of three equations with three unknowns: a , ρ and ω . However, Eq.(27) can also be obtained by rearranging (25) and (26). In this way, of these three equations, two are independent, what makes the system insoluble.

What one makes, then, is to assume a specific value for the EoS ω and solve the Friedmann equations for a and ρ . Usually, one takes $\omega = 1/3$ and $\omega = 0$ for radiation and matter components, respectively. On the other hand, the accelerated stages, namely inflationary and dark energy eras, can be described by an exotic fluid of negative EoS, specifically with $\omega < -1/3$ [38].

Anyhow, so far there is no analytical approach in SM able to describe a unique functional form for $\omega = \omega(t)$ that describes the different stages of the Universe as well as the respective transitions.

In the present article, from the application of the first order formalism in Section III, we were able to construct the cosmological scenarios of Section IV. In the next paragraphs we will check the reliability of these scenarios and its relation with the values of the free parameters of the models. For now, it is important to remark that we obtained a complete set of exact solutions for the studied models, which display double and single-kink configurations, as we can see from Figs.1 and 2. In fact, such models are very important in applications that include Bloch branes, Skyrmions, Yang-Mills, Q-balls, oscillons and various superstring-motivated theories.

Let us start by analysing the Hubble factor (Fig.3). Qualitatively, all curves are in accord with SM in the sense that $H \propto t_H^{-1}$, with t_H being the Hubble time. Still in agreement with SM, H is restricted to positive values ($H < 0$ would imply a contracting universe).

One might wonder the reason why $H(t) \rightarrow \text{constant}$ for

high values of time. In order to interpret such a behavior of $H(t)$, one should keep in mind that $H = \dot{a}/a$ and that an accelerated expansion might be described by an exponential of t as $a \sim e^{H_0 t}$, with H_0 a constant (in this case, the present value of $H(t)$). Such a solution for $a(t)$ yields a constant behavior for $H(t)$, therefore $H = \text{constant}$ in Fig.3 for high values of time stands for the recent regime of accelerated expansion the Universe is passing through.

Fig.4 shows the time evolution of the EoS parameter of Model I and II. As described some paragraphs above, in SM one assumes to know the values of this parameter for different eras of the Universe, but there is no mechanism capable to predict the functional form of $\omega(t)$. Here, on the other hand, we have obtained from the model, a solution to $\omega(t)$ and this is describing the whole evolution of the Universe dynamics as we will argue in the following.

Firstly we note that ω starts in negative values, specifically, when $t \sim 0$, $\omega \sim -1$, independently of the model. This is in agreement with some constraints that were recently put to the inflationary EoS [49–51]. Note that this behavior is reflected in the curves of $H(t)$ for small values of time ($t \rightarrow 0 \Rightarrow H(t) \rightarrow \text{constant}$).

After describing inflation, ω tends to positive values. Note that for the values chosen in Fig.4 for the constants of the model, ω has its maximum value around 1/3. Indeed, that is the maximum value the EoS of the Universe should assume according to SM, as carefully described above. In this way, our models predict a smooth transition between inflationary and radiation eras, which is not trivial to be obtained, even with alternative models (check Refs.[52, 53] for some content about the so-called “graceful exit”).

The EoS parameter, then, starts to decrease, eventually re-assuming negative values. Note that for higher values of time, $\omega \rightarrow -1$, which is related with the cosmic acceleration and is in accordance with the present value of the EoS, obtained via observation of fluctua-

tions of the cosmic microwave background temperature ($\omega = -1.073^{+0.090}_{-0.089}$) [4].

Let us now check Fig.5, in which the acceleration parameter is depicted. For small values of time, $\bar{q} > 0$, which stands for an accelerated stage of the Universe expansion. According to the argumentations above, we can conclude that the behavior of $\bar{q}$ for small values of is nothing but the description of the inflationary era.

As time passes by, $\bar{q}$ tends to negative values, which describe a decelerated expansion of the Universe. As also argued above, we know the dynamics of the Universe expansion changes with time. After inflationary era, the Universe expanded in a decelerated way, during radiation and matter epochs. Those are described by the period of time in which $\bar{q} < 0$ in Fig.7.

After that, $\bar{q}$ turns back to positive values, passes through its current value [4], in accordance with the present cosmic acceleration, and tends to 1, asymptotically predicting, in this way, a de Sitter-like universe in the far future. It is worth, here, remarking that the prediction of an asymptotically de Sitter universe is also reflected in Fig.3, for the Hubble parameter, since the de Sitter universe is described by a constant Hubble factor.

In this way, our multi-quintessence models have shown to be able to provide a complete cosmological scenario. The different dynamical stages of the Universe were described, as well as the transitions among them, in a continuous form.

Acknowledgments

RACC thanks CAPES for financial support. PHRSM thanks FAPESP, Grant No. 2015/08476-0, for financial support.

-
- [1] A. G. Riess et al., *Astron. J.* **116**, 1009 (1998).
 - [2] G. Perlmutter et al., *Astrophys. J.* **517**, 565 (1999).
 - [3] S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989).
 - [4] G. Hinshaw et al., *Astrophys. J.* **208**, 19 (2013).
 - [5] T. P. Sotiriou and V. Faraoni, *Rev. Mod. Phys.* **82**, 451 (2010).
 - [6] A. de Felice and S. Tsujikawa, *Liv. Rev. Rel.* **13**, 3 (2010).
 - [7] S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **74**, 086005 (2006).
 - [8] S. Nojiri and S. D. Odintsov, *Phys. Rep.* **505**, 59 (2011).
 - [9] T. Harko et al., *Phys. Rev. D* **84**, 024020 (2011).
 - [10] P.H.R.S. Moraes, *Eur. Phys. J. C* **75**, 168 (2015).
 - [11] P.H.R.S. Moraes, *Astrophys. Space Sci.* **352**, 273 (2014).
 - [12] P.H.R.S. Moraes and R.A.C. Correa, *Astrophys. Space Sci.* **361**, 91 (2016).
 - [13] P.H.R.S. Moraes, *Int. J. Theor. Phys.* **55**, 1307 (2016).
 - [14] P.H.R.S. Moraes and J.R.L. Santos, *Eur. Phys. J. C* **76**, 60 (2016).
 - [15] V. Faraoni, *Phys. Rev. D* **62**, 023504 (2000).
 - [16] E. Elizalde et al., *Phys. Rev. D* **77**, 106005 (2008).
 - [17] X.-F. Zhang et al., *Mod. Phys. Lett. A* **21**, 231 (2006).
 - [18] M. Khurshudyan et al., *Int. J. Theor. Phys.* **53**, 2370 (2014).
 - [19] M. C. Bento et al., *Phys. Rev. D* **65**, 067301 (2002).
 - [20] P. H. R. S. Moraes and J.R.L. Santos, *Phys. Rev. D* **89**, 083516 (2014).
 - [21] D. Bazeia, C. B. Gomes, L. Losano, and R. Menezes, *Phys. Lett. B* **633**, 415 (2006).
 - [22] A. de Souza Dutra, G. P. de Brito, and J. M. Hoff, *Phys. Rev. D* **91**, 086016 (2015).
 - [23] R. Rajaraman, *Solitons and Instantons* (North-Holland, Amsterdam, 1982).
 - [24] T. Vachaspati, *Kinks and Domain Walls: An Introduction to Classical and Quantum Solitons* (Cambridge University Press, Cambridge, England, 2006).
 - [25] D. Bazeia, J. R. S. Nascimento, R. F. Ribeiro, and D.

- Toledo, J. Phys. A **30**, 8157 (1997).
- [26] D. Bazeia and F.A. Brito, Phys. Rev. D **61**, 105019 (2000).
 - [27] M. B. Voloshin, Phys. Rev. D **57**, 1266 (1997).
 - [28] M. Shifman, Phys. Rev. D **57**, 2358 (1997).
 - [29] M. A. Shifman and M.B. Voloshin, Phys. Rev. D **57**, 2590 (1998).
 - [30] E. Ventura, A.M. Simas, and D. Bazeia, Chem. Phys. Lett. **320**, 587 (2000).
 - [31] R. A. C. Correa, A. de Souza Dutra, M. B. Hott, Class. Quant. Grav. **28**, 155012 (2011).
 - [32] A. de Souza Dutra and R. A. C. Correa, Phys.Rev. D **83**, 105007 (2011).
 - [33] R. A. C. Correa, R. da Rocha, and A. de Souza Dutra, Annals Phys. **359**, 198 (2015).
 - [34] A. de Souza Dutra, M. Hott, and F.A. Barone, Phys. Rev. D **74**, 085030 (2006).
 - [35] M. Gleiser, N. Stamatopoulos, Phys. Lett. B **713**, 304 (2012).
 - [36] R. A. C. Correa, A. de Souza Dutra, and M. Gleiser, Phys. Lett. B **737**, 388 (2014).
 - [37] A. de Souza Dutra, Phys. Lett. B **626**, 249 (2005).
 - [38] B. Ryden, *Introduction to Cosmology* (Addison Wesley, San Francisco, USA, 2003).
 - [39] S. Weinberg, *Cosmology* (Oxford University Press, New York, USA, 2008).
 - [40] S. Nojiri and S.D. Odintsov, Phys. Rev. D **78**, 046006 (2008).
 - [41] W. Hu and I. Sawicki, Phys. Rev. D **76**, 064004 (2007).
 - [42] E. Elizalde et al., Phys. Rev. D **70**, 043539 (2004).
 - [43] L. Amendola, Phys. Rev. D **62**, 043511 (2000).
 - [44] S. Das and N. Banerjee, Gen. Rel. Grav. **38**, 785 (2006).
 - [45] F. Wilczek, Nuclear Phys. B Proceedings Suppl. **134**, 3 (2004).
 - [46] S. Capozziello and M. Francaviglia, Gen. Rel. Grav. **40**, 357 (2008).
 - [47] A.H. Guth, Phys. Rev. D **23**, 347 (1981).
 - [48] J.D. Barrow and M.S. Turner, Nature **292**, 35 (1981).
 - [49] K. Bamba et al., Class. Quant. Grav. **30**, 015008 (2013).
 - [50] L. Ackerman et al., JCAP **05**, 024 (2011).
 - [51] R. Aldrovani et al., Eur. Phys. J. C **58**, 483 (2008).
 - [52] J.A.S. Lima et al., MNRAS **431**, 923 (2013).
 - [53] E.L.D. Perico et al., Phys. Rev. D **88**, 063531 (2013)